arXiv:1504.01305v1 [math.CV] 6 Apr 2015

THE PSEUDOANALYTIC EXTENSIONS FOR SOME SPACES OF ANALYTIC FUNCTIONS

GUANLONG BAO, HASI WULAN AND FANGQIN YE

ABSTRACT. Using the Cauchy-Riemann operator, we characterize $\mathcal{Q}_K$ spaces, Besov spaces and analytic Morrey spaces in terms of pseudoanalytic extensions of primitive functions. Our results are also true on some classical Banach spaces, such as the Bloch space, $BMOA$ and the Dirichlet space.

Keywords: the Cauchy-Riemann operator; pseudoanalytic extension; primitive functions; $\mathcal{Q}_K$ spaces; Besov spaces; Morrey spaces.

1. INTRODUCTION

Let $\mathbb{D}$ be the unit disk in the complex plane $\mathbb{C}$. Denote by $H(\mathbb{D})$ the space of functions analytic in $\mathbb{D}$. The Green function in the unit disk with singularity at $a \in \mathbb{D}$ is given by

$$g(a,z) = \log \frac{1}{|\sigma_a(z)|},\ z \in \mathbb{D}.$$

Here

$$\sigma_a(z) = \frac{a-z}{1-\overline{a}z},$$

is a Möbius transformation of $\mathbb{D}$.

Throughout this paper, we assume that $K : [0,\infty) \to [0,\infty)$ is a right-continuous and increasing function. A function $f \in H(\mathbb{D})$ belongs to the space $\mathcal{Q}_K$ if

$$\|f\|^2_{\mathcal{Q}_K} = \sup_{a\in\mathbb{D}} \int_{\mathbb{D}} |f'(z)|^2 K\left(g(a,z)\right) dA(z) < \infty,$$

where $dA(z)$ is the Lebesgue measure in $\mathbb{D}$. By [14, Theorem 2.1], $f \in \mathcal{Q}_K$ if and only if

$$\sup_{a\in\mathbb{D}} \int_{\mathbb{D}} |f'(z)|^2 K\left(1-|\sigma_a(z)|^2\right) dA(z) < \infty.$$

See [14] and [15] for more results of $\mathcal{Q}_K$ spaces. If $K(t) = t^p$, $0 \le p < \infty$, then the space $\mathcal{Q}_K$ gives the space $\mathcal{Q}_p$ (see [5, 29, 30]). In particular, $\mathcal{Q}_0$ is the Dirichlet space $\mathcal{D}$; $\mathcal{Q}_1 = BMOA$, the space of bounded mean oscillation (see [6, 17]); by [3], for all $p \in (1,\infty)$, the spaces $\mathcal{Q}_p$ are the same and equal to the Bloch space $\mathcal{B}$ which consists of all functions $f \in H(\mathbb{D})$ with

$$\|f\|_{\mathcal{B}} = \sup_{z\in\mathbb{D}} (1-|z|^2)|f'(z)| < \infty.$$

2000 *Mathematics Subject Classification.* 30H25, 46E15.

The authors are supported by NSF of China (No. 11371234).

For $0<p<\infty$, the Hardy space H^p consists of all functions $f\in H(\mathbb{D})$ with

$$\|f\|_{H^p}^p=\sup_{0<r<1}\frac{1}{2\pi}\int_0^{2\pi}|f(re^{i\theta})|^p d\theta<\infty.$$

It is well-known that if $f\in H^p$, then its nontangential limit $f(e^{i\theta})$ exists almost everywhere (see [9, 16]).

In this paper, we need the Cauchy-Riemann operator

$$\overline{\partial}=\frac{\partial}{\partial\overline{z}}=\frac{1}{2}\left(\frac{\partial}{\partial x}+i\frac{\partial}{\partial y}\right),\ z=x+iy.$$

Dyn'kin [13] said that the so-called pseudoanalytic extension is a method of extension with a given estimate of the Cauchy-Riemann operator. Dyn'kin [13] characterized many classical smoothness spaces applying the pseudoanalytic extension method. Note that

$$\bigcup_{0<p<1}\mathcal{Q}_p\subseteq BMOA\subseteq\bigcap_{0<q<\infty}H^q.$$

Dyakonov and Girela [12] obtained an interesting characterization of $\mathcal{Q}_p$ spaces in terms of pseudoanalytic extension as follows.

Theorem A. *If $0<p<1$ and $f\in\bigcap_{0<q<\infty}H^q$, then the following conditions are equivalent.*

(i) $f\in\mathcal{Q}_p$.

(ii)

$$\sup_{a\in\mathbb{D}}\int_{\mathbb{D}}|f'(z)|^2\left(\frac{1}{|\sigma_a(z)|^2}-1\right)^p dA(z)<\infty.$$

(iii) There exists a function $\widetilde{f}\in C^1(\mathbb{C}\setminus\overline{\mathbb{D}})$ satisfying

$$\widetilde{f}(z)=O(1),\ \textit{as}\ z\to\infty,$$

$$\lim_{r\to1^+}\widetilde{f}(re^{i\theta})=f(e^{i\theta}),\ \textit{a.e. and in}\ L^q([-\pi,\ \pi])\ \textit{for all}\ q\in[1,\infty),$$

and

$$\sup_{a\in\mathbb{D}}\int_{\mathbb{C}\setminus\overline{\mathbb{D}}}|\overline{\partial}\widetilde{f}(z)|^2\left(|\sigma_a(z)|^2-1\right)^p dA(z)<\infty.$$

The proof of the above theorem in [12] involved the Calderón-Zygmund operators and the Muckenhoupt weights (see [19]). For $a\in\mathbb{D}$ and $0<p<1$, let

$$U_a(z)=\left|1-\frac{1}{|\sigma_a(z)|^2}\right|^p=\frac{(1-|a|^2)^p||z|^2-1|^p}{|z-a|^{2p}},\ z\in\mathbb{C}.$$

Dyakonov and Girela [12] showed that U_a is a Muckenhoupt weight with

$$\sup_Q\left[\frac{1}{|Q|}\int_Q U_a(z)dA(z)\right]\left[\frac{1}{|Q|}\int_Q(U_a(z))^{-1}dA(z)\right]<\infty$$

for all $a\in\mathbb{D}$. Here Q ranges over the disks in $\mathbb{C}$ and $|Q|$ denotes the area of Q. The above estimate of U_a with $a=0$ was also used to establish the $\mathcal{Q}_p$ corona theorem (see [28, Theorem 3.1]). Applying Theorem A, Dyakonov and Girela [12] also gave some nice properties of $\mathcal{Q}_p$ spaces.

Motivated by Theorem A, it is nature to ask some questions as follows.

Question 1. Can we obtain pseudoanalytic extensions for all $\mathcal{Q}_p$, $0 \le p < \infty$?

Question 2. Let n be a positive integer. Does there exist a pseudoanalytic extension characterization of $\mathcal{Q}_p$ spaces such that the extension function is $O(z^n)$ as $z \to \infty$?

Question 3. Can we give characterizations for some spaces of analytic functions such as $\mathcal{Q}_K$ spaces, Besov spaces and analytic Morrey spaces in terms of pseudoanalytic extensions?

Throughout this article, denote by F the primitive function of $f \in H(\mathbb{D})$; that is

$$F(z) = \int_0^z f(w)dw, \ z \in \mathbb{D}.$$

Related to the above questions, in this paper, we characterize some analytic function spaces on $\mathbb{D}$ by pseudoanalytic extension of primitive functions. Our method, without using Calderón-Zygmund operators and Muckenhoupt weights, can be applied to some spaces, such as $\mathcal{Q}_K$ spaces, Besov spaces, analytic Morrey spaces and so on. Our results are also new for $\mathcal{Q}_p$ spaces.

In this paper, the symbol $A \approx B$ means that $A \lesssim B \lesssim A$. We say that $A \lesssim B$ if there exists a constant C such that $A \le CB$.

2. $\mathcal{Q}_K$ SPACES AND PSEUDOANALYTIC EXTENSION

In this section, we need two more constraints on K as follows.

$$\int_0^1 \frac{\varphi_K(s)}{s} ds < \infty \tag{2.1}$$

and

$$\int_1^\infty \frac{\varphi_K(s)}{s^{1+q}} ds < \infty, \quad 0 < q < 2, \tag{2.2}$$

where

$$\varphi_K(s) = \sup_{0 \le t \le 1} \frac{K(st)}{K(t)}, \quad 0 < s < \infty.$$

Under conditions (2.1) and (2.2), $\mathcal{Q}_K$ spaces have been studied extensively (see [24, 25, 26]). From now on we always assume the function K satisfying the double condition, namely $K(2t) \approx K(t)$ for all $t \in (0, 1)$.

A very useful tool in the study of $\mathcal{Q}_K$ spaces is K-Carleson measure. Let $\ell(I)$ be the length of an arc I of the unit circle $\partial\mathbb{D}$. Define the Carleson box by

$$S_G(I) = \begin{cases} \{r\zeta \in G : 1 - \frac{\ell(I)}{2\pi} < r < 1, \zeta \in I\}, & G = \mathbb{D}, \\ \{r\zeta \in G : 1 < r < 1 + \ell(I), \zeta \in I\}, & G = \mathbb{C} \setminus \overline{\mathbb{D}}. \end{cases}$$

Following [15] and [24], a positive Borel measure μ on $G = \mathbb{D}$ or $G = \mathbb{C} \setminus \overline{\mathbb{D}}$ is said to be a K-Carleson measure if

$$\sup_{I \subset \partial\mathbb{D}} \int_{S_G(I)} K\left(\frac{|1 - |z||}{\ell(I)}\right) d\mu(z) < \infty.$$

By [14], we know that all $\mathcal{Q}_K$ spaces are subsets of the Bloch space. Then the primitive function F of a $\mathcal{Q}_K$ function f must be in the Hardy space H^2 since the Bloch functions' Taylor coefficients is bounded (see [1]). Therefore, the primitive function F has its nontangential limit $F(e^{i\theta})$ almost everywhere on the unit circle.

The following is the main result of this section.

Theorem 2.1. *Suppose that K satisfies (2.1) and (2.2). Let $f \in H(\mathbb{D})$ with its primitive function $F \in H^2$ and let $n \geq 2$ be an integer. Then the following conditions are equivalent.*

(i) $f \in \mathcal{Q}_K$.

(ii) There exists a function $\widetilde{F_n} \in C^1(\mathbb{C} \setminus \overline{\mathbb{D}})$ satisfying

$$\lim_{r \to 1^+} \widetilde{F_n}(re^{i\theta}) = F(e^{i\theta}) \ a.e. \ \theta \in [0, 2\pi], \tag{2.3}$$

$$\widetilde{F_n}(z) = O(z^n), \text{ as } z \to \infty, \tag{2.4}$$

$$\overline{\partial}\widetilde{F_n}(z) = O(z^{n-2}), \text{ as } z \to \infty, \tag{2.5}$$

and

$$\sup_{a \in \mathbb{D}} \int_{\mathbb{C}\setminus\overline{\mathbb{D}}} \frac{|\overline{\partial}\widetilde{F_n}(z)|^2}{(|z|^n - 1)^2} K\left(1 - \frac{1}{|\sigma_a(z)|^2}\right) dA(z) < \infty. \tag{2.6}$$

(iii) There exists a function $\widetilde{F_n} \in C^1(\mathbb{C}\setminus\overline{\mathbb{D}})$ satisfying (2.3), (2.4), (2.5) and $|\overline{\partial}\widetilde{F_n}(z)|^2/(|z|^n - 1)^2$ is a K-Carleson measure on $\mathbb{C} \setminus \overline{\mathbb{D}}$.

Before embarking into the proof of Theorem 2.1, we state some lemmas below.

Lemma 2.2. *Suppose that K satisfies (2.1). Let μ be a positive Borel measure on $\mathbb{D}$. Then the following conditions are equivalent.*

(i) μ is a K-Carleson measure on $\mathbb{D}$.

(ii)

$$\sup_{a \in \mathbb{D}} \int_{\mathbb{D}} K(1 - |\sigma_a(z)|^2) d\mu(z) < \infty.$$

(iii)

$$\sup_{a \in \mathbb{D}} \int_{\mathbb{D}} K\left(\frac{1}{|\sigma_a(z)|^2} - 1\right) d\mu(z) < \infty.$$

Proof. Note that $(i) \Leftrightarrow (ii)$ was proved in [15]. To show the equivalence of (ii) and (iii), let

$$K_1(t) = K\left(\frac{t}{1-t}\right), \ 0 < t < \frac{1}{2}.$$

By the monotonicity of K, $K_1(s)$ is also increasing for $s \in (0, 1)$. Moreover,

$$K(t) \leq K\left(\frac{t}{1-t}\right) \leq K(2t), \ 0 < t < \frac{1}{2}.$$

Since $K(2t) \approx K(t)$ for $t \in (0, 1)$, $K_1(t) \approx K(t)$ for $t \in (0, 1/2)$. Thus, the proof is complete. □

Wulan and Zhu [27] proved that if K satisfies (2.1) and n is a positive integer, then $f \in \mathcal{Q}_K$ if and only if

$$\sup_{a\in\mathbb{D}} \int_{\mathbb{D}} |f^{(n)}(z)|^2(1-|z|^2)^{2n-2}K\left(1-|\sigma_a(z)|^2\right)dA(z) < \infty. \tag{2.7}$$

Combining this with Lemma 2.2, we obtain immediately a characterization of $\mathcal{Q}_K$ spaces as follows. If K satisfies (2.1) and n is a positive integer, then $f \in \mathcal{Q}_K$ if and only if

$$\sup_{a\in\mathbb{D}} \int_{\mathbb{D}} |f^{(n)}(z)|^2(1-|z|^2)^{2n-2}K\left(\frac{1}{|\sigma_a(z)|^2}-1\right)dA(z) < \infty.$$

For the case of μ on $\mathbb{C}\setminus\overline{\mathbb{D}}$, we give a similar description of K-Carleson measure as follows.

Lemma 2.3. *Suppose that K satisfies (2.1). Let μ be a positive Borel measure on $\mathbb{C}\setminus\overline{\mathbb{D}}$. Then the following conditions are equivalent.*

(i) *μ is a K-Carleson measure on $\mathbb{C}\setminus\overline{\mathbb{D}}$.*

(ii)

$$\sup_{a\in\mathbb{D}} \int_{1<|z|<1+2\pi} K\left(|\sigma_a(z)|^2-1\right)d\mu(z) < \infty.$$

(iii)

$$\sup_{a\in\mathbb{D}} \int_{1<|z|<1+2\pi} K\left(1-\frac{1}{|\sigma_a(z)|^2}\right)d\mu(z) < \infty.$$

Proof. $(i) \Rightarrow (ii)$. Fix $a = re^{i\theta} \in \mathbb{D}$. Let I be the arc of center $e^{i\theta}$ and $\ell(I) = \frac{2\pi(1-|a|)}{2\pi|a|+1}$. Then for any $z \in S_{\mathbb{C}\setminus\overline{\mathbb{D}}}(I)$, one gets

$$|1-\bar{a}z| \geq 1-(1+\ell(I))|a| = \frac{1}{2\pi}\ell(I)$$

and

$$\begin{aligned} |1-\bar{a}z| &\leq |a|\left|e^{i\left(\theta+\frac{\ell(I)}{2}\right)}-\frac{1}{\bar{a}}\right| \\ &= r\left|e^{i\frac{\ell(I)}{2}}-\frac{1}{r}\right| = \left[(1-r)^2+4r\left(\sin\frac{\ell(I)}{4}\right)^2\right]^{\frac{1}{2}} \\ &\leq \left[\left(\frac{2\pi r+1}{2\pi}\ell(I)\right)^2+4r\left(\frac{\ell(I)}{4}\right)^2\right]^{\frac{1}{2}} \leq 2\ell(I). \end{aligned}$$

Hence

$$|1-\bar{a}z| \approx 1-|a| \approx \ell(I), \quad z \in S_{\mathbb{C}\setminus\overline{\mathbb{D}}}(I).$$

Set

$$S_n = \left\{r\zeta \in \mathbb{C}\setminus\overline{\mathbb{D}} : 1 < r < 1+\ell(2^nI), \zeta \in 2^nI\right\},$$

where $2^n I$ denotes the arc with the same center as I and with length $2^n\ell(I)$. Then

$$\begin{aligned}
&\int_{1<|z|<1+2\pi} K\left(|\sigma_a(z)|^2-1\right)d\mu(z)\\
\leq\ & \int_{S_{\mathbb{C}\setminus\overline{\mathbb{D}}}(I)} K\left(|\sigma_a(z)|^2-1\right)d\mu(z)+\sum_{n=1}^{\infty}\int_{S_n\setminus S_{n-1}} K\left(|\sigma_a(z)|^2-1\right)d\mu(z)\\
\lesssim\ & \int_{S_{\mathbb{C}\setminus\overline{\mathbb{D}}}(I)} K\left(\frac{|z|-1}{\ell(I)}\right)d\mu(z)+\sum_{n=1}^{\infty}\int_{S_n\setminus S_{n-1}} K\left(\frac{|z|-1}{2^{2n}\ell(I)}\right)d\mu(z)\\
\lesssim\ & \int_{S_{\mathbb{C}\setminus\overline{\mathbb{D}}}(I)} K\left(\frac{|z|-1}{\ell(I)}\right)d\mu(z)+\sum_{n=1}^{\infty}\sup_{z\in S_n}\frac{K\left(\frac{|z|-1}{2^{2n}\ell(I)}\right)}{K\left(\frac{|z|-1}{2^{n}\ell(I)}\right)}\int_{S_n} K\left(\frac{|z|-1}{\ell(2^nI)}\right)d\mu(z)\\
\lesssim\ & \int_{S_{\mathbb{C}\setminus\overline{\mathbb{D}}}(I)} K\left(\frac{|z|-1}{\ell(I)}\right)d\mu(z)+\sum_{n=1}^{\infty}\varphi_K(2^{-n})\int_{S_n} K\left(\frac{|z|-1}{\ell(2^nI)}\right)d\mu(z).
\end{aligned}$$

Since $d\mu$ is a K-Carleson measure and

$$\sum_{n=1}^{\infty}\varphi_K(2^{-n})\approx\int_0^1\varphi_K(s)\frac{ds}{s}<\infty,$$

we obtain that

$$\sup_{a\in\mathbb{D}}\int_{1<|z|<1+2\pi} K\left(|\sigma_a(z)|^2-1\right)d\mu(z)<\infty.$$

$(ii)\Rightarrow(iii)$. It follows from that

$$K\left(1-\frac{1}{|\sigma_a(z)|^2}\right)\leq K\left(|\sigma_a(z)|^2-1\right)$$

holds for all $a\in\mathbb{D}$ and $1<|z|<1+2\pi$.

$(iii)\Rightarrow(i)$. For given a subarc $I\in\partial\mathbb{D}$, set $e^{i\theta}$ the midpoint of I and set

$$a=\frac{2\pi-\ell(I)}{2\pi(\ell(I)+1)}e^{i\theta}.$$

Then for any $z\in S_{\mathbb{C}\setminus\overline{\mathbb{D}}}(I)$, we have

$$0<|\sigma_a(z)|^2-1\leq\frac{\frac{1}{\pi}(2\pi|a|+1)(2+\ell(I))\left(\ell(I)\right)^2}{\left(\frac{1}{2\pi}\ell(I)\right)^2}\leq 8\pi(2\pi+1)(\pi+1).$$

Let

$$K_2(t)=K\left(\frac{t}{1+t}\right),\ 0<t<8\pi(2\pi+1)(\pi+1).$$

By the monotonicity of K, $K_2(s)$ is also increasing for $s\in(0,1)$. Moreover,

$$K\left(\frac{t}{8\pi(2\pi+1)(\pi+1)+1}\right)\leq K_2(t)\leq K(t),\ 0<t<8\pi(2\pi+1)(\pi+1).$$

Since $K(2t)\approx K(t)$ for $t\in(0,1)$, $K_2(t)\approx K(t)$ for $t\in(0,8\pi(2\pi+1)(\pi+1))$. Thus,

$$\begin{aligned}\int_{S_{\mathbb{C}\setminus\overline{\mathbb{D}}}(I)} K\left(\frac{|z|-1}{\ell(I)}\right)d\mu(z) &\approx \int_{S_{\mathbb{C}\setminus\overline{\mathbb{D}}}(I)} K\left(|\sigma_a(z)|^2-1\right)d\mu(z)\\ &\approx \int_{S_{\mathbb{C}\setminus\overline{\mathbb{D}}}(I)} K\left(\frac{|\sigma_a(z)|^2-1}{|\sigma_a(z)|^2}\right)d\mu(z)\\ &\lesssim \int_{1<|z|<1+2\pi} K\left(1-\frac{1}{|\sigma_a(z)|^2}\right)d\mu(z).\end{aligned}$$

Hence, (i) follows. □

From [15] and [25], if K satisfies (2.1) and (2.2), then there exists a small enough constant c, $0<c<q/2$, such that $K(t)/t^c$ is increasing and $K(t)/t^{q-c}$ is decreasing in (0, 1). To prove Theorem 2.1, we also need the following estimate.

Lemma 2.4. *Let (2.1) and (2.2) hold for K. If $s<\min(1+c,2-q+c)$ and $2s+r-4\geq 0$, then*

$$\int_{\mathbb{D}}\frac{K\left(1-|\sigma_a(w)|^2\right)}{(1-|w|^2)^s|1-\overline{w}z|^r}dA(w)\lesssim\frac{K\left(1-|\sigma_a(z)|^2\right)}{(1-|z|^2)^{s+r-2}}$$

for all $a,z\in\mathbb{D}$. Here c is a small enough positive constant depending only on (2.1) and (2.2).

Proof. For fixed a, $z\in\mathbb{D}$, let $\lambda=\sigma_z(a)$. Then

$$|\sigma_a(w)|=|\sigma_\lambda\circ\sigma_z(w)|.$$

Note that $2s+r-4\geq 0$. Checking the proof of Lemma 2.1 in [7], one gets

$$\begin{aligned}&\int_{\mathbb{D}}\frac{K\left(1-|\sigma_a(w)|^2\right)}{(1-|w|^2)^s|1-\overline{w}z|^r}dA(w)\\ \lesssim\ &\frac{K\left(1-|\lambda|^2\right)}{(1-|z|^2)^{s+r-2}}\int_{\mathbb{D}}\varphi_K\left(\frac{1-|u|^2}{|1-\overline{\lambda}u|^2}\right)\frac{1}{(1-|u|^2)^s}dA(u).\end{aligned}$$

Since K satisfies (2.1) and (2.2), there exists a small enough positive constant c depending only on (2.1) and (2.2) (see [15, 25]), such that

$$\varphi_K(t)\lesssim t^c,\ 0<t\leq 1$$

and

$$\varphi_K(t)\lesssim t^{q-c},\ t\geq 1.$$

Thus,

$$\begin{aligned}&\int_{\mathbb{D}}\varphi_K\left(\frac{1-|u|^2}{|1-\overline{\lambda}u|^2}\right)\frac{1}{(1-|u|^2)^s}dA(u)\\ \lesssim\ &\int_{\mathbb{D}}\frac{(1-|u|^2)^{c-s}}{|1-\overline{\lambda}u|^{2c}}dA(u)+\int_{\mathbb{D}}\frac{(1-|u|^2)^{q-c-s}}{|1-\overline{\lambda}u|^{2q-2c}}dA(u).\end{aligned}$$

For $s<1+c<q+1-c$ and $s<2-q+c$, using Lemma 3.10 in [35], we get

$$\int_{\mathbb{D}}\frac{(1-|u|^2)^{c-s}}{|1-\overline{\lambda}u|^{2c}}dA(u)+\int_{\mathbb{D}}\frac{(1-|u|^2)^{q-c-s}}{|1-\overline{\lambda}u|^{2q-2c}}dA(u)\lesssim 1.$$

Thus,

$$\int_{\mathbb{D}} \frac{K\left(1-|\sigma_a(w)|^2\right)}{(1-|w|^2)^s|1-\overline{w}z|^r}dA(w) \lesssim \frac{K\left(1-|\lambda|^2\right)}{(1-|z|^2)^{s+r-2}} \approx \frac{K\left(1-|\sigma_a(z)|^2\right)}{(1-|z|^2)^{s+r-2}}.$$

□

Now we are ready to finish the proof of Theorem 2.1.

$(i) \Rightarrow (ii)$. From now on, we write

$$w^* = 1/\overline{w},\ w \in \mathbb{C} \setminus \{0\}.$$

Let $f \in \mathcal{Q}_K$ and

$$F(z) = \int_0^z f(\zeta)d\zeta,\ z \in \mathbb{D}.$$

Set

$$\widetilde{F_n}(z) = \sum_{i=0}^{n} \frac{(-1)^i}{i!}(z^*-z)^i F^{(i)}(z^*),\ z \in \mathbb{C} \setminus \overline{\mathbb{D}}.$$

Clearly, $\widetilde{F_n} \in C^1(\mathbb{C} \setminus \overline{\mathbb{D}})$ satisfying

$$\lim_{r\to 1^+} \widetilde{F_n}(re^{i\theta}) = F(e^{i\theta})\ \ a.e.\ \theta \in [0, 2\pi],$$

and

$$\widetilde{F_n}(z) = O(z^n),\ \text{as}\ \ z \to \infty.$$

Note that

$$\overline{\partial}\widetilde{F_n}(z) = \frac{(-1)^{n+1}}{n!}(z^*-z)^n(z^*)^2F^{(n+1)}(z^*).$$

Then

$$\overline{\partial}\widetilde{F_n}(z) = O(z^{n-2}),\ \text{as}\ \ z \to \infty.$$

Making the change of variable $z = \zeta^*$, we deduce that

$$\begin{aligned} &\sup_{a\in\mathbb{D}} \int_{\mathbb{C}\setminus\overline{\mathbb{D}}} \frac{\left|\overline{\partial}\widetilde{F_n}(z)\right|^2}{(|z|^n-1)^2} K\left(1-\frac{1}{|\sigma_a(z)|^2}\right) dA(z) \\ =\ & \frac{1}{(n!)^2}\sup_{a\in\mathbb{D}} \int_{\mathbb{C}\setminus\overline{\mathbb{D}}} \frac{|z^*-z|^{2n}|z^*|^4\left|f^{(n)}(z^*)\right|^2}{(|z|^n-1)^2} K\left(1-\frac{1}{|\sigma_a(z)|^2}\right) dA(z) \\ \approx\ & \sup_{a\in\mathbb{D}} \int_{\mathbb{D}} \left|f^{(n)}(\zeta)\right|^2 (1-|\zeta|)^{2n-2} K\left(1-|\sigma_a(\zeta)|^2\right) dA(\zeta). \end{aligned}$$

Combining this with (2.7), we get the desired result.

$(ii) \Rightarrow (i)$. Assume that (ii) holds. Let $z \in \mathbb{D}$ and $R > 1$. Following a technique in [10] or [12] and in view of (2.3), we employ the Cauchy-Green formula to the function that equals F in $\mathbb{D}$ and $\widetilde{F_n}$ in $\mathbb{C} \setminus \overline{\mathbb{D}}$. Then

$$F(z) = \frac{1}{2\pi i}\int_{|\zeta|=1} \frac{F(\zeta)}{\zeta - z}d\zeta = \frac{1}{2\pi i}\int_{|\zeta|=R} \frac{\widetilde{F_n}(\zeta)}{\zeta - z}d\zeta - \frac{1}{\pi}\int_{1<|\zeta|<R} \frac{\overline{\partial}\widetilde{F_n}(\zeta)}{\zeta - z}dA(\zeta). \tag{2.8}$$

By (2.4),

$$\int_{|\zeta|=R} \frac{\widetilde{F_n}(\zeta)}{(\zeta - z)^{n+2}} d\zeta \to 0, \text{ as } R \to \infty.$$

By (2.5), we know

$$\left| \int_{\mathbb{C}\backslash\overline{\mathbb{D}}} \frac{\overline{\partial}\widetilde{F_n}(\zeta)}{(\zeta - z)^{n+2}} dA(\zeta) \right| < \infty.$$

These together with (2.8) give

$$F^{(n+1)}(z) = -\frac{(n+1)!}{\pi} \int_{\mathbb{C}\backslash\overline{\mathbb{D}}} \frac{\overline{\partial}\widetilde{F_n}(\zeta)}{(\zeta - z)^{n+2}} dA(\zeta).$$

By the Hölder inequality, the change of variable and the well known estimates in Zhu's book [35, Lemma 3.10], we obtain

$$\begin{aligned} \left|F^{(n+1)}(z)\right|^2 &\lesssim \int_{\mathbb{C}\backslash\overline{\mathbb{D}}} \frac{1}{|\zeta - z|^4} dA(\zeta) \int_{\mathbb{C}\backslash\overline{\mathbb{D}}} \frac{|\overline{\partial}\widetilde{F_n}(\zeta)|^2}{|\zeta - z|^{2n}} dA(\zeta) \\ &\approx \int_{\mathbb{D}} \frac{1}{|1 - \overline{w}z|^4} dA(w) \int_{\mathbb{C}\backslash\overline{\mathbb{D}}} \frac{|\overline{\partial}\widetilde{F_n}(\zeta)|^2}{|\zeta - z|^{2n}} dA(\zeta) \\ &\approx \frac{1}{(1-|z|^2)^2} \int_{\mathbb{C}\backslash\overline{\mathbb{D}}} \frac{|\overline{\partial}\widetilde{F_n}(\zeta)|^2}{|\zeta - z|^{2n}} dA(\zeta). \end{aligned}$$

Using Lemma 2.4, $n \geq 2$ and (2.6), we see that

$$\begin{aligned} & \sup_{a\in\mathbb{D}} \int_{\mathbb{D}} (1-|z|^2)^{2n-2} \left|F^{(n+1)}(z)\right|^2 K\left(1 - |\sigma_a(z)|^2\right) dA(z) \\ \lesssim\ & \sup_{a\in\mathbb{D}} \int_{\mathbb{D}} \int_{\mathbb{C}\backslash\overline{\mathbb{D}}} \frac{|\overline{\partial}\widetilde{F_n}(\zeta)|^2}{|\zeta - z|^{2n}} dA(\zeta) (1-|z|^2)^{2n-4} K\left(1 - |\sigma_a(z)|^2\right) dA(z) \\ \lesssim\ & \sup_{a\in\mathbb{D}} \int_{\mathbb{D}} \int_{\mathbb{D}} \frac{K\left(1 - |\sigma_a(z)|^2\right)}{|1 - \overline{w}z|^4} dA(z) |\overline{\partial}\widetilde{F_n}(w^*)|^2 |w|^{2n-4} dA(w) \\ \lesssim\ & \sup_{a\in\mathbb{D}} \int_{\mathbb{D}} \frac{|\overline{\partial}\widetilde{F_n}(w^*)|^2 |w|^{2n-4}}{(1-|w|^2)^2} K\left(1 - |\sigma_a(w)|^2\right) dA(w) \\ \approx\ & \sup_{a\in\mathbb{D}} \int_{\mathbb{C}\backslash\overline{\mathbb{D}}} \frac{|\overline{\partial}\widetilde{F_n}(\zeta)|^2}{(|\zeta|^n - 1)^2} K\left(1 - \frac{1}{|\sigma_a(\zeta)|^2}\right) dA(\zeta) < \infty. \end{aligned}$$

Note that $F'(z) = f(z)$ for $z \in \mathbb{D}$. By (2.7), we get $f \in \mathcal{Q}_K$.

$(ii) \Leftrightarrow (iii)$. If

$$\overline{\partial}\widetilde{F_n}(z) = O(z^{n-2}), \text{ as } z \to \infty,$$

then

$$\sup_{a\in\mathbb{D}} \int_{|z|\geq 1+2\pi} \frac{|\overline{\partial}\widetilde{F_n}(z)|^2}{(|z|^n - 1)^2} K\left(1 - \frac{1}{|\sigma_a(z)|^2}\right) dA(z) < \infty.$$

This together with Lemma 2.3, we see that condition (ii) is equivalent to condition (iii). We finish the proof. □

Remark. In Theorem 2.1, by the particular choice of K, we obtain the corresponding characterizations of $\mathcal{Q}_p$ for $0 < p < 1$, $BMOA$ and the Bloch space.

3. BESOV SPACES, MORREY SPACES AND PSEUDOANALYTIC EXTENSION

In this section, we describe Besov spaces and Morrey spaces by pseudoanalytic extension of primitive functions.

For $1 < p < \infty$, the Besov space B_p is the space of analytic functions f in $\mathbb{D}$ such that

$$\|f\|_{B_p}^p = \int_{\mathbb{D}} |f'(z)|^p (1-|z|^2)^{p-2} dA(z) < \infty. \tag{3.1}$$

In particular, if $p = 2$, then B_p is the Dirichlet space $\mathcal{D}$. Note that (3.1) does not hold for $p = 1$. The Besov space B_1 consists of all analytic functions f on $\mathbb{D}$ which have a representation as

$$f(z) = \sum_{k=1}^{\infty} c_k \sigma_{a_k}(z), \ a_k \in \mathbb{D} \ \text{ and } \sum_{k=1}^{\infty} |c_k| < \infty.$$

Form [2], $f \in B_1$ if and only if

$$\int_{\mathbb{D}} |f''(z)| dA(z) < \infty.$$

Let $1 \le p < \infty$ and $n \ge 2$ be an integer. It is well known that $f \in B_p$ if and only if

$$\int_{\mathbb{D}} |f^{(n)}(z)|^p (1-|z|^2)^{np-2} dA(z) < \infty. \tag{3.2}$$

See [11, 34, 35] for more results of Besov spaces.

Note that all Besov spaces are subsets of the Bloch space. Thus, if $f \in B_p$, then its primitive function F belongs to the Hardy space H^2. Now we give a pseudoanalytic extension characterization of Besov spaces as follows.

Theorem 3.1. *Let $f \in H(\mathbb{D})$ with its primitive function $F \in H^2$ and let $n \ge 2$ be an integer. If $1 \le p < \infty$, then the following conditions are equivalent.*

(i) $f \in B_p$.

(ii) There exists a function $\widetilde{F_n} \in C^1(\mathbb{C} \setminus \overline{\mathbb{D}})$ satisfying

$$\lim_{r \to 1^+} \widetilde{F_n}(re^{i\theta}) = F(e^{i\theta}) \ a.e. \ \theta \in [0, 2\pi], \tag{3.3}$$

$$\widetilde{F_n}(z) = O(z^n), \ \textit{as} \ \ z \to \infty, \tag{3.4}$$

$$\overline{\partial}\widetilde{F_n}(z) = O(z^{n-2}), \ \textit{as} \ \ z \to \infty, \tag{3.5}$$

and

$$\int_{\mathbb{C}\setminus\overline{\mathbb{D}}} \frac{|\overline{\partial}\widetilde{F_n}(z)|^p}{\left(|z|^{\frac{np}{2}-p+2} - 1\right)^2} dA(z) < \infty. \tag{3.6}$$

Proof. $(i) \Rightarrow (ii)$. Let $f \in B_p$. Set

$$\widetilde{F_n}(z) = \sum_{i=0}^{n} \frac{(-1)^i}{i!}(z^* - z)^i F^{(i)}(z^*),\ z \in \mathbb{C} \setminus \overline{\mathbb{D}}.$$

It is easy to check that $\widetilde{F_n} \in C^1(\mathbb{C} \setminus \overline{\mathbb{D}})$ and $\widetilde{F_n}$ satisfies (3.3), (3.4) and (3.5). Furthermore, by the following estimates

$$\begin{aligned}
& \int_{\mathbb{C}\setminus\overline{\mathbb{D}}} \frac{|\overline{\partial}\widetilde{F_n}(z)|^p}{\left(|z|^{\frac{np}{2}-p+2}-1\right)^2} dA(z) \\
= \ & \frac{1}{(n!)^p} \int_{\mathbb{C}\setminus\overline{\mathbb{D}}} \frac{|z^*-z|^{np}|z^*|^{2p}\left|f^{(n)}(z^*)\right|^p}{\left(|z|^{\frac{np}{2}-p+2}-1\right)^2} dA(z) \\
\approx \ & \int_{\mathbb{D}} \frac{(1-|w|^2)^{np}\left|f^{(n)}(w)\right|^p}{\left(1-|w|^{\frac{np}{2}-p+2}\right)^2} dA(w) \\
\approx \ & \int_{\mathbb{D}} (1-|w|)^{np-2}|f^{(n)}(w)|^p dA(w)
\end{aligned}$$

and (3.2), we know that $\widetilde{F_n}$ also satisfies (3.6).

$(ii) \Rightarrow (i)$. Checking the proof of Theorem 2.1, one gets

$$F^{(n+1)}(z) = -\frac{(n+1)!}{\pi} \int_{\mathbb{C}\setminus\overline{\mathbb{D}}} \frac{\overline{\partial}\widetilde{F_n}(\zeta)}{(\zeta-z)^{n+2}} dA(\zeta),\ z \in \mathbb{D}.$$

Applying the Hölder inequality and [35, Lemma 3.10], we obtain

$$\begin{aligned}
\left|F^{(n+1)}(z)\right|^p \ & \lesssim \left(\int_{\mathbb{C}\setminus\overline{\mathbb{D}}} \frac{1}{|\zeta-z|^4} dA(\zeta)\right)^{p-1} \int_{\mathbb{C}\setminus\overline{\mathbb{D}}} \frac{|\overline{\partial}\widetilde{F_n}(\zeta)|^p}{|\zeta-z|^{np-2p+4}} dA(\zeta) \\
& \approx \left(\int_{\mathbb{D}} \frac{1}{|1-\overline{w}z|^4} dA(w)\right)^{p-1} \int_{\mathbb{C}\setminus\overline{\mathbb{D}}} \frac{|\overline{\partial}\widetilde{F_n}(\zeta)|^p}{|\zeta-z|^{np-2p+4}} dA(\zeta) \\
& \approx \frac{1}{(1-|z|^2)^{2p-2}} \int_{\mathbb{C}\setminus\overline{\mathbb{D}}} \frac{|\overline{\partial}\widetilde{F_n}(\zeta)|^p}{|\zeta-z|^{np-2p+4}} dA(\zeta).
\end{aligned}$$

Note that $n \geq 2$. Using [35, Lemma 3.10] again, we deduce that

$$\begin{aligned}
&\int_{\mathbb{D}} (1-|z|^2)^{np-2} \left|F^{(n+1)}(z)\right|^p dA(z) \\
\lesssim\ & \int_{\mathbb{D}} \int_{\mathbb{C}\backslash\overline{\mathbb{D}}} \frac{|\overline{\partial}\widetilde{F_n}(\zeta)|^p}{|\zeta - z|^{np-2p+4}} dA(\zeta)(1-|z|^2)^{np-2p} dA(z) \\
\approx\ & \int_{\mathbb{D}} \int_{\mathbb{D}} \frac{|\overline{\partial}\widetilde{F_n}(w^*)|^p |w|^{np-2p}}{|1-\overline{w}z|^{np-2p+4}} dA(w)(1-|z|^2)^{np-2p} dA(z) \\
\approx\ & \int_{\mathbb{D}} \int_{\mathbb{D}} \frac{(1-|z|^2)^{np-2p}}{|1-\overline{w}z|^{np-2p+4}} dA(z) |\overline{\partial}\widetilde{F_n}(w^*)|^p |w|^{np-2p} dA(w) \\
\lesssim\ & \int_{\mathbb{D}} \left(1-|w|^2\right)^{-2} |\overline{\partial}\widetilde{F_n}(w^*)|^p |w|^{np-2p} dA(w) \\
\approx\ & \int_{\mathbb{D}} \left(1-|w|^{\frac{np}{2}-p+2}\right)^{-2} |\overline{\partial}\widetilde{F_n}(w^*)|^p |w|^{np-2p} dA(w) \\
\approx\ & \int_{\mathbb{C}\backslash\overline{\mathbb{D}}} \frac{|\overline{\partial}\widetilde{F_n}(\zeta)|^p}{(|\zeta|^{\frac{np}{2}-p+2}-1)^2} dA(\zeta).
\end{aligned}$$

Since $F'(z) = f(z)$ for $z \in \mathbb{D}$, by (3.2) and (3.6), we obtain $f \in B_p$. This finishes the proof. $\square$

For $0 < \lambda \leq 1$, the analytic Morrey space $\mathcal{L}^{2,\lambda}$ is the set of all functions $f \in H^2$ satisfying

$$\|f\|_{\mathcal{L}^{2,\lambda}} = \sup_{I \subseteq \partial\mathbb{D}} \left(\frac{1}{\ell(I)^\lambda} \int_I |f(\zeta) - f_I|^2 |d\zeta| \right)^{1/2} < \infty,$$

where $\ell(I)$ denotes the length of I and

$$f_I = \frac{1}{\ell(I)} \int_I f(\zeta) |d\zeta|.$$

Clearly, $\mathcal{L}^{2,1}$ is the space $BMOA$. By [30, p. 54], $f \in \mathcal{L}^{2,\lambda}$ if and only if

$$\sup_{a \in \mathbb{D}} (1-|a|^2)^{1-\lambda} \int_{\mathbb{D}} |f'(z)|^2 (1-|\sigma_a(z)|^2) dA(z) < \infty.$$

Let $n \geq 1$ be an integer. Using a similar statement in [4], we see that $f \in \mathcal{L}^{2,\lambda}$ if and only if

$$\sup_{a \in \mathbb{D}} (1-|a|^2)^{1-\lambda} \int_{\mathbb{D}} |f^{(n)}(z)|^2 (1-|z|^2)^{2n-2} (1-|\sigma_a(z)|^2) dA(z) < \infty. \tag{3.7}$$

Recently, the interest in analytic Morrey space has grown rapidly. See [8, 18, 20, 21, 23, 31, 32] for more results of $\mathcal{L}^{2,\lambda}$ spaces.

Note that if $f \in \mathcal{L}^{2,\lambda}$, then its primitive function F belongs to H^2. Applying (3.7), we get the following characterization of analytic Morrey space. The proof is similar to Theorem 2.1, we omit it.

Theorem 3.2. *Let $f \in H(\mathbb{D})$ with its primitive function $F \in H^2$ and let $n \geq 2$ be an integer. If $0 < \lambda \leq 1$, then the following conditions are equivalent.*

(i) $f \in \mathcal{L}^{2,\lambda}$.

(ii) There exists a function $\widetilde{F_n} \in C^1(\mathbb{C} \setminus \overline{\mathbb{D}})$ *satisfying*

$$\lim_{r\to 1^+} \widetilde{F_n}(re^{i\theta}) = F(e^{i\theta}) \; a.e. \; \theta \in [0, 2\pi],$$

$$\widetilde{F_n}(z) = O(z^n), \text{ as } z \to \infty,$$

$$\overline{\partial}\widetilde{F_n}(z) = O(z^{n-2}), \text{ as } z \to \infty,$$

and

$$\sup_{a\in\mathbb{D}}(1-|a|^2)^{1-\lambda}\int_{\mathbb{C}\setminus\overline{\mathbb{D}}} \frac{|\overline{\partial}\widetilde{F_n}(z)|^2}{(|z|^n-1)^2}\left(1-\frac{1}{|\sigma_a(z)|^2}\right)dA(z) < \infty.$$

Remark. When the paper is complete, we found that Wei and Shen obtained the special case of our Theorem 2.1 for $K(t) = t$ and $n = 2$. However, our proof in this paper and those of [22] are different.

REFERENCES

[1] J. Anderson, J. Clunie and Ch. Pommerenke, On Bloch functions and normal functions, *J. Reine Angew. Math.*, **270**(1974), 12-37.

[2] J. Arazy, S. Fisher and J. Peetre, Möbius invariant function spaces, *J. Reine Angew. Math.*, **363** (1985), 110-145.

[3] R. Aulaskari and P. Lappan, Criteria for an analytic function to be Bloch and a harmonic or meromorphic function to be normal, *Complex analysis and its applications, Pitman Res. Notes in Math., 305, Longman Sci. Tec., Harlow,* 1994, 136-146.

[4] R. Aulaskari, M. Nowak and R. Zhao, The nth derivative characterisation of Möbius invariant Dirichlet space, *Bull. Austral. Math. Soc.*, **58**(1998), 43-56.

[5] R. Aulaskari, J. Xiao and R. Zhao, On subspaces and subsets of $BMOA$ and UBC, *Analysis.*, **15**(1995), 101-121.

[6] A. Baernstein, Analytic functions of Bounded Mean Oscillation. In: Aspects of Con- temporary Complex Analysis, Editors: D. A. Brannan and J. G. Clunie., Academic Press, London, New York (1980), pp. 3-36.

[7] G. Bao, Z. Lou, R. Qian and H. Wulan, Improving multipliers and zero sets in $\mathcal{Q}_K$ spaces, *Collect. Math.*, doi: 10.1007/s13348-014-0113-z.

[8] C. Cascante, J. Fabrega and J. Ortega, The corona theorem in weighted Hardy and Morrey spaces, *Ann. Scuola Norm. Super. Pisa Cl. Sci.*, doi: 10.2422/2036-2145.201202-006.

[9] P. Duren, Theory of H^p Spaces, Academic Press, New York, 1970.

[10] K. Dyakonov, Equivalent norms on Lipschitz-type spaces of holomorphic functions, *Acta Math.*, **178**(1997), 143-167.

[11] K. Dyakonov, Besov spaces and outer functions, *Michigan Math. J.*, **45**(1998), 143-157.

[12] K. Dyakonov and D. Girela, On $\mathcal{Q}_p$ spaces and pseudoanalytic extension, *Ann. Acad. Sci. Fenn. Math.*, **25**(2000), 477-486.

[13] E. Dyn'kin, The pseudoanalytic extension, *J. Anal. Math.* , **60**(1993), 45-70.

[14] M. Essén and H. Wulan, On analytic and meromorphic function and spaces of $\mathcal{Q}_K$-type, *Illionis. J. Math.*, **46**(2002), 1233-1258.

[15] M. Essén, H. Wulan and J. Xiao, Several function-theoretic characterizations of Möbius invariant $\mathcal{Q}_K$ spaces, *J. Funct. Anal.*, **230**(2006), 78-115.

[16] J. Garnett, Bounded Analytic Functions, Springer, New York, 2007.

[17] D. Girela, Analytic functions of bounded mean oscillation. In: Complex Function Spaces, Mekrijärvi 1999 Editor: R. Aulaskari. Univ. Joensuu Dept. Math. Rep. Ser. 4, Univ. Joensuu, Joensuu (2001) pp. 61-170.

[18] P. Li, J. Liu and Z. Lou, Integral operators on analytic Morrey spaces, *Sci. China Math.*, **57**(2014), 1961-1974.

[19] E. Stein, Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals, Princeton University Press, Princeton, NJ, 1993.

[20] J. Wang and J. Xiao, Two predualities and three operators over analytic campanato spaces, arXiv:1402.4377.

[21] J. Wang and J. Xiao, Holomorphic campanato spaces on the unit ball, arXiv:1405.6192.

[22] H. Wei and Y. Shen, On the tangent space to the BMO-Teichmüller space, *J. Math. Anal. Appl.*, **419**(2014), 715-726.

[23] Z. Wu and C. Xie, $\mathcal{Q}$ spaces and Morrey spaces, *J. Funct. Anal.*, **201**(2003), 282-297.

[24] H. Wulan and F. Ye, Universal Teichmüller space and $\mathcal{Q}_K$ spaces, *Ann. Acad. Sci. Fenn. Math.*, **39**(2014), 691-709.

[25] H. Wulan and J. Zhou, Decomposition theorems for $\mathcal{Q}_K$ spaces and applications, *Forum Math.*, **26**(2014), 467-495.

[26] H. Wulan and J. Zhou, $\mathcal{Q}_K$ and Morrey type spaces, *Ann. Acad. Sci. Fenn. Math.*, **38**(2013), 193-207.

[27] H. Wulan, K. Zhu, $\mathcal{Q}_K$ spaces via higher order derivatives, *Rocky Mountain J. Math.*, **38**(2008), 329-350.

[28] J. Xiao, The $\mathcal{Q}_p$ corona theorem, *Pacific J. Math.*, **194**(2000), 491-509.

[29] J. Xiao, Holomorphic $\mathcal{Q}$ Classes, Springer, LNM 1767, Berlin, 2001.

[30] J. Xiao, Geometric $\mathcal{Q}_p$ functions, Birkhäuser Verlag, Basel-Boston-Berlin, 2006.

[31] J. Xiao and W. Xu, Composition operators between analytic Campanato spaces, *J. Geom. Anal.*, **24**(2014), 649-666.

[32] J. Xiao and C. Yuan, Analytic Campanato spaces and their compositions, arXiv:1303.5032.

[33] R. Zhao, Distances from Bloch functions to some Möbius invariant spaces, *Ann. Acad. Sci. Fenn. Math.*, **33**(2008), 303-313.

[34] K. Zhu, Analytic Besov spaces, *J. Math. Anal. Appl.*, **157** (1991), 318-336.

[35] K. Zhu, Operator Theory in Function Spaces, American Mathematical Society, Providence, RI, 2007.

DEPARTMENT OF MATHEMATICS, SHANTOU UNIVERSITY, SHANTOU, GUANGDONG, 515063, CHINA
E-mail address: `glbaoah@163.com (G. Bao)`
E-mail address: `wulan@stu.edu.cn (H. Wulan)`
E-mail address: `yefqah@163.com (F. Ye)`